

## Data Sheets on Quarantine Pests

*Trioza erytrae***IDENTITY**

**Name:** *Trioza erytrae* (Del Guercio)

**Synonyms:** *Spanioza erytrae* Del Guercio  
*Trioza merwei* Pettey

**Taxonomic position:** Insecta: Hemiptera: Homoptera: Psyllidae

**Common names:** Citrus psylla (English)  
Citrusblattfloh (German)

**Bayer computer code:** TRIZER

**EPPO A1 list:** No. 46

**EU Annex designation:** II/A1

**HOSTS**

*T. erytrae* is confined to Rutaceae, occurring on wild hosts (*Clausena anisata*, *Vespris undulata*) as well as on *Citrus*, especially lemons (*C. limon*) and limes (*C. aurantiifolia*). Within the EPPO region the host species are generally confined to countries surrounding the Mediterranean.

**GEOGRAPHICAL DISTRIBUTION**

The distribution of *T. erytrae* is wider than that of citrus greening bacterium, the major pathogen which it transmits (EPPO/CABI, 1996a), since it occurs in Cameroon, Congo, Malawi, Rwanda, St. Helena, Sudan, Tanzania, Uganda and Zambia where the bacterium has not been recorded.

**EPPO region:** Portugal (Madeira only; isolated outbreaks first found in 1994).

**Asia:** Saudi Arabia, Yemen.

**Africa:** Cameroon, Comoros, Ethiopia, Kenya, Madagascar, Malawi, Mauritius, Réunion, Rwanda, South Africa, St. Helena, Sudan, Swaziland, Tanzania, Uganda, Zaire, Zambia, Zimbabwe. The record in Congo which appeared in the previous version of this data sheet (EPPO/CABI, 1992) is a mistake; it correctly refers to Zaire.

**EU:** Present.

**Distribution map:** See CIE (1967, No. 234).

**BIOLOGY**

A bibliography of *T. erytrae* up to 1987 has been compiled by Van den Berg & Fletcher (1988) and a general review has been presented by Van den Berg (1990). *T. erytrae* has a temperature sensitivity similar to that of the African form of citrus greening bacterium (Schwarz & Green, 1970; Catling, 1973). It is very sensitive to extremes of hot, dry weather (the eggs and first instar nymphs being particularly vulnerable). It is favoured in cool, moist areas over 500-600 m, where citrus growth flushes tend to be prolonged. Green

& Catling (1971) have used maximum saturation deficit as an accurate predictor of the geographical distribution of *T. erythrae*.

Sex ratios fluctuate in the field, but females always predominate. There is a pre-oviposition period of 3-7 days, but this is considerably extended in the absence of young foliage; longevity is also prolonged under such conditions. Mating occurs 2 to 4 times a day and eggs may be laid immediately. Eggs are supplied with a sharp point which is driven through the leaf epidermis and is thought to be responsible for maintaining a favourable internal water relationship. Females remain fertile for 11-16 days in the absence of males, and maximum egg production occurs towards the middle of their life span, which normally lasts 17-50 days; up to 2000 eggs may be laid per female. There is an incubation period of 6-15 days and nymphal development (five instars) takes 17-43 days, both periods being inversely related to mean temperature and directly related to nutritional value of the leaves. The temperature threshold for nymphal development is, apparently, around 10-12°C. There is no diapause. Van den Berg *et al.* (1990) have recently studied the daily activities and habits of adults, and egg hatching and moulting, in *T. erythrae*, while Van den Berg *et al.* (1991a) have studied mating, fertility and oviposition.

*T. erythrae* transmits the African form of citrus greening bacterium under natural conditions in Africa and Yemen (McClellan & Oberholzer, 1965). It has been shown experimentally that *T. erythrae* is also able to transmit the Asian form (Massonie *et al.*, 1976). In Mauritius and Réunion, where both forms occur, *T. erythrae* probably transmits both.

## DETECTION AND IDENTIFICATION

### Symptoms

*T. erythrae* severely distorts leaves, which are stunted and galled, and appear dusted with fecal pellets.

### Morphology

#### Eggs

Orange, cylindrical, with a sharp point anteriorly; laid on leaf margins of young, actively growing foliage.

#### Nymph

Dorso-ventrally compressed and varying in colour from yellow, olive-green to dark-grey; has a marginal fringe of white, waxy filaments; largely sedentary; forms distinct colonies and settles on the underside of young leaves, where, after a few days' feeding, it produces distinctive cup-shaped, open galls.

#### Adult

Winged, pale and delicate initially, later becoming light-brown. Males are smaller than females and have a blunt tip to the abdomen, the latter ending in a sharp point in females. When feeding, adults take up a distinctive stance, with the abdomen raised at an angle of about 35° to the feeding surface.

## MEANS OF MOVEMENT AND DISPERSAL

*T. erythrae* is only liable to be locally spread by natural dispersal, up to distances of 1.5 km (Van den Berg & Deacon, 1988). Citrus material (budwood, grafted trees, rootstock seedlings) from infected areas can carry eggs and/or nymphs over longer distances. Such 5th or 6th-instar nymphs, as well as the adults born from nymphs, are capable of transmitting the greening agent to citrus. Entry on citrus fruit is extremely unlikely.

## PEST SIGNIFICANCE

### Economic impact

The main economic importance of *T. erytreae* is as the vector of the very serious citrus disease caused by citrus greening bacterium (EPPO/CABI, 1996a). Heavy infestations with *T. erytreae* also cause severe leaf distortion and development of conspicuous pits on the leaf surface.

### Control

Insecticides such as dimethoate can be used against *T. erytreae*, for which there is an active monitoring programme in orchards in South Africa. *T. erytreae* enters orchards from indigenous hosts in surrounding vegetation (Van den Berg *et al.*, 1991b), so it is recommended to remove them. In Réunion, *T. erytreae* has been successfully controlled by the introduction of a parasite, *Tamarixia dryi*, from South Africa (Aubert *et al.*, 1980). In South Africa, numerous predators occur but have not been found to reduce populations to economically acceptable levels (Van den Berg *et al.*, 1987).

### Phytosanitary risk

Like the other vector of citrus greening (*Diaphorina citri*; EPPO/CABI, 1996b), *T. erytreae* is listed as an A1 quarantine pest by EPPO (OEPP/EPPO, 1988) and is also a quarantine pest for CPPC and OIRSA. It could probably establish and spread in Mediterranean countries without difficulty. Besides its role in citrus greening, this psyllid has significant damage potential in itself. Though biological control may be possible, there is no guarantee that it could keep populations to a sufficiently low level to prevent transmission of greening.

## PHYTOSANITARY MEASURES

EPPO recommends (OEPP/EPPO, 1990) that importation of plants for planting and cut branches of citrus from countries where citrus greening bacterium or either of its vectors occur should be prohibited. It is possible to fumigate citrus budwood material against *T. erytreae* (FAO, 1983).

## BIBLIOGRAPHY

- Aubert, B.; Bové, J.M.; Etienne, J. (1980) La lutte contre la maladie du greening des agrumes à l'île de la Réunion. Résultats et perspectives. *Fruits* **35**, 605-624.
- Catling, H.D. (1973) Notes on the biology of the South African citrus psylla *Trioza erytreae*. *Journal of the Entomological Society of South Africa* **36**, 299-306.
- CIE (1967) *Distribution Maps of Pests, Series A* No. 234. CAB International, Wallingford, UK.
- EPPO/CABI (1992) *Trioza erytreae*. In: *Quarantine pests for Europe* (Ed. by Smith, I.M.; McNamara, D.G.; Scott, P.R.; Harris, K.M.). CAB International, Wallingford, UK.
- EPPO/CABI (1996a) Citrus greening bacterium. In: *Quarantine pests for Europe*. 2nd edition (Ed. by Smith, I.M.; McNamara, D.G.; Scott, P.R.; Holderness, M.). CAB INTERNATIONAL, Wallingford, UK.
- EPPO/CABI (1996b) *Diaphorina citri*. In: *Quarantine pests for Europe*. 2nd edition (Ed. by Smith, I.M.; McNamara, D.G.; Scott, P.R.; Holderness, M.). CAB INTERNATIONAL, Wallingford, UK.
- FAO (1983) International plant quarantine treatment manual. *FAO Plant Production and Protection Paper* No. 50. FAO, Rome, Italy.
- Green, G.C.; Catling, H.D. (1971) Weather-induced mortality of the citrus psylla *Trioza erytreae*, a vector of greening virus, in some citrus-producing areas of South Africa. *Agricultural Meteorology* **8**, 305-317.

- Massonie, G.; Garnier, M.; Bové, J.M. (1976) Transmission of Indian citrus decline by *Trioza erytreae*, the vector of South African greening. In: *Proceedings of the 7th Conference of the International Organization of Citrus Virologists* (Ed. by Calavan, E.C.), pp. 18-20. University of California, Riverside, USA.
- McClellan, A.P.D.; Oberholzer, P.C.J. (1965) Citrus psylla, a vector of the greening disease of sweet orange. *South African Journal of Agricultural Science* **8**, 297-298.
- OEPP/EPPO (1988) Data sheets on quarantine organisms No. 151, Citrus greening bacterium and its vectors *Diaphorina citri* & *Trioza erytreae*. *Bulletin OEPP/EPPO Bulletin* **18**, 497-507.
- OEPP/EPPO (1990) Specific quarantine requirements. *EPPO Technical Documents* No. 1008.
- Schwarz, R.E.; Green, G.C. (1970) Citrus greening and the citrus psyllid *Trioza erytreae*, a temperature-dependent agent-vector complex. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* **79**, 490-493.
- Van den Berg, M.A. (1990) The citrus psylla, *Trioza erytreae*: a review. *Agriculture, Ecosystems and Environment* **30**, 171-194.
- Van den Berg, M.A.; Deacon, V.E. (1988) Dispersal of the citrus psylla, *Trioza erytreae*, in the absence of its host plants. *Phytophylactica* **20**, 361-368.
- Van den Berg, M.A.; Fletcher, C.D. (1988) A bibliography of the citrus psylla, *Trioza erytreae*, up to 1987. *Phytoparasitica* **16**, 47-61.
- Van den Berg, M.A.; Deacon, V.E.; Fourie, C.J.; Anderson, S.H. (1987) Predators of the citrus psylla, *Trioza erytreae*, in the Lowveld and Rustenburg areas of Transvaal. *Phytophylactica* **19**, 285-289.
- Van den Berg, M.A.; Deacon, V.E.; Jager, K. de (1990) Ecology of the citrus psylla, *Trioza erytreae*. 1. Daily activities and habits of adults. 2. Egg hatching and moulting. *Phytophylactica* **22**, 323-328, 329-333.
- Van den Berg, M.A.; Deacon, V.E.; Thomas, C.D. (1991a) Ecology of the citrus psylla, *Trioza erytreae*. 3. Mating, fertility and oviposition. *Phytophylactica* **23**, 195-200.
- Van den Berg, M.A.; Deacon, V.E.; Steenekamp, P.J. (1991b) Dispersal within and between citrus orchards and native hosts, and nymphal mortality of citrus psylla, *Trioza erytreae*. *Agriculture, Ecosystems and Environment* **35**, 297-309.