

# Validating and using new methods for detecting plant diseases

**Glyn Jones** 

Principal Economist, Fera

**Roy Macarthur** 

Statistician, Fera

### Validating and using a new detection method

- Validation?
- Mathematical odyssey
- Example: screening symptomatic plants for phytophthora ramorum using a \*new LFD\* before a more expensive established test
- Don't wait until you have a detection method to validate it.

### Validating a new method?

- My new method; what is it good for?
- Estimate real world outcomes with and without the detection technology using well designed validation data
- So, for a rapid on-site LFD test...

#### In a world...

...without LFDs

Test plants by looking for symptoms; send symptomatic plants to be tested in a lab

• ... with LFDs

Test plants by looking for symptoms; test plants with LFDs, maybe based on symptoms; send plants to be tested in a lab depending on symptoms and LFD


What are costs and outcomes in each world

#### Validation data?

 Test infected and uninfected plants as assessed by a gold standard method; report false negative and false positive rates.


#### Validation data

• Test some symptomatic and non-symptomatic plants by all methods and also get an estimate of the proportion symptomatic.  $N_1$ 




Each set of plants at a lower level is either all of the plants from a higher level, or a random sample from the higher level

## Validation data: cost of an inspection plan



Without LFDs  $Cost = inspection + p_a \times Lab$ 


### Validation data: cost of an inspection plan



Without LFDs  $Cost = inspection + p_a \times Lab$ 

Maybe cheaper with LFD


Validation data: performance of an inspection plan



Without LFDs: Infected=  $(1 - p_a)(p_b p_e + (1 - p_b)p_d)$ 

With LFDs: Infected=  $(1 - p_a)(p_b p_e + (1 - p_b)p_d) + p_a(1 - p_c)p_f$ 

Validation data: performance of an inspection plan




Without LFDs: Infected=  $(1 - p_a)(p_b p_e + (1 - p_b)p_d)$ 


With LFDs: Infected=  $(1 - p_a)(p_b p_e + (1 - p_b)p_d) + p_a(1 - p_c)p_f$ 

Maybe more infected plants slip through

#### Observations




### Uncertainty associated with estimates





### Uncertainty associated with derived estimates:


Proportion infected in plants that pass a an inspection plan


Infected= 
$$(1 - p_a)(p_b p_e + (1 - p_b)p_d) + p_a(1 - p_c)p_f$$














### Uncertainty associated with derived estimates:


Proportion infected in plants that pass a an inspection plan


Infected= 
$$(1 - p_a)(p_b p_e + (1 - p_b)p_d) + p_a(1 - p_c)p_f$$














# Is using the LFD as a screening test in inspection useful?

Test plan	Cost per plant inspected(£) Central estimate and 95% C.I			Proportion of infected plants among those that "pass" (%) Central estimate and 95% C.I			
DO NOTHING	0	0	0	4.3	2.1	11.5	
Send symptomatic to Lab	1.38	1.13	1.68	3.8	1.6	11.1	
Test symptomatic with LFD; send +ve to Lab	0.93	0.73	1.18	3.8	1.6	11.1	
Send symptomatic and asymptomatic that give a +ve LFD to Lab	19.2	17.9	26.3	2.6	0.79	8.8	

# What if the LFD worked twice as well at half the cost?

Test plan	Cost per plant inspected(£)  Central estimate and 95% C.I			Proportion of infected plants among those that "pass" (%) Central estimate and 95% C.I			
DO NOTHING	0	0	0	4.3	2.1	11.5	
Send symptomatic to Lab	1.38	1.13	1.68	3.8	1.6	11.1	
Test symptomatic with LFD; send +ve to Lab	0.92	0.72	1.17	3.8	1.6	11.0	
Send symptomatic and asymptomatic that give a +ve LFD to Lab	16.8	14.6	24.8	1.3	0.31	6.9	

# Is using the LFD as a screening test during surveillance against a possible outbreak useful?

Scenario	Illustrative Infection prevalence at first detection(%) Estimate 95% CI			Relative Prevalence at first detection Estimate 95% CI		
Send symptomatic to Lab	0.121	0.0593	0.336	1	1	1
Test symptomatic with LFD; send +ve to Lab	0.0909	0.0458	0.253	0.750	0.660	0.873
Send symptomatic and asymptomatic that give a +ve LFD to Lab	0.469	0.288	1.75	3.87	1.55	10.5
Test symptomatic with cheaper improved LFD; send +ve to Lab	(0.0853)	0.0439	0.244	0.735	0.648	0.852

Based on rule of thumb described by Parnell et al (2015), modified for probability of detecting an infected plant

#### What if....?

What if we had a device with similar performance to the LFD, but allowing the examination of a large number of plants in a on-site bulk test, or maybe by multispectral examination?

### Rapid bulk screening for inspection

Test plan	Cost per plant inspected(£) Central estimate and 95% C.I			Proportion of infected plants among those that "pass" Central estimate and 95% C.I			
DO NOTHING	0	0	0	4.3	2.1	11.5	
Send symptomatic to Lab	1.38	1.13	1.68	3.8	1.6	11.1	
Test symptomatic with screen; send +ve to Lab	0.77	0.59	0.99	3.9	1.7	11.1	
Send symptomatic and asymptomatic that give a +ve screen to Lab	3.00	1.76	10.1	2.6	0.079	8.8	

# Rapid bulk screening for surveillance against a new outbreak

Scenario	Illustrative* Infection prevalence at first detection(%) Estimate 95% CI			Relative Prevalence at first detection Estimate 95% CI		
Send symptomatic to Lab	0.121	0.0593	0.336	1	1	1
Test symptomatic with LFD; send +ve to Lab	0.0909	0.0458	0.253	0.750	0.660	0873
Test symptomatic with cheaper improved LFD; send +ve to Lab	0.0853	0.0439	0.244	0.735	0.648	0.852
Test symptomatic with bulk screen; send +ve to Lab	0.0749	0.0377	0.208	0.618	0.528	0.732
Test asymptomatic with bulk screen; send +ve to Lab along with symptomatic	0.0735	0.0516	0.300	0.606	0.498	1.74

#### Reflections

The validation was based on 160 tests; by careful design we can get a lot of information from those tests beyond the false positive and negative rates of a single method.

Using the framework to model potential technologies as well as different inspection plans can help focus plans.