

EXTRAPOLATION TABLE for EFFECTIVENESS of INSECTICIDES

► PESTS ON STONE FRUIT

INTRODUCTION

The table provides detailed lists of acceptable extrapolations organized by crop groups, for regulatory authorities and applicants, in the context of the registration of plant protection products for minor uses. The table should be used in conjunction with the EPPO Standard PP1/257(1) - *Efficacy and crop safety extrapolations for minor uses*. It is important to ensure that expert judgment and regulatory experience are employed when using these tables. EPPO excludes liability as to the reliability of the information provided through these tables.

The scope for extrapolation may be extended as data and experience with a certain plant protection products increases. The applicant should always provide appropriate justification and information to support the proposed extrapolation. For example, comparability of target biology may be a relevant factor, either in extrapolating to other target species or for the same target onto another crop. For crops, factors such as comparable growth habit, structure etc. should be considered.

TABLE FORMAT

The main pest species for the crop group are listed in Column 1 (although this is not exhaustive), and the pest group to which they belong is specified in Column 2. Companies may choose if they wish to provide data only for individual named species, which would then appear individually listed on the label. But underlined species have been identified as key major targets and as such it is advisable to generate data on these. Furthermore, data on these species then allow a claim to be made for the whole pest group (as specified in Column 2), if required. If a claim for the whole pest group is required but there is no underlined species, then data must be generated on all listed species.

Column 3 indicates the key indicator crop(s) for the crop group. In some instances this may be only one specified crop. In other cases, when separated by an 'or', the company may choose from a range of alternatives within the group. Data generated on crops in Column 3 may be used to extrapolate to all crops listed in Column 4. However, it is preferable to have data on several of the crops within the crop group, but data on the indicator crop should be available. In specific circumstances data from crops outside of the crop group highlighted by an asterisk in column 5 can replace the need for any data on the indicator crop in column 3.

Column 5 identifies whether relevant data on crops outside the crop group, against the same target, may help to reduce the amount of required data on the indicator crop. It may be possible for a direct extrapolation without the need for data on the indicator crop (marked with an asterisk

(*). However, this is dependent on the extent of available data and similarity of crop/target biology. The company should provide an appropriate reasoned case when wanting to use data from crops outside the crop group.

Column 6 gives examples of acceptable extrapolations for a particular pest claim onto other minor use crops. This is not a comprehensive list. Whether extrapolation may be direct (no data, marked with an asterisk (*)), or require additional supporting data on the minor use crop, will again be dependent on the extent and relevance of the existing database and companies should provide an appropriate reasoned case. If the crop is considered to be a major crop in some countries then it may not be appropriate to include in this column, and further data would be required. Companies will need to justify the status of the major crop/minor use.

EXAMPLE OF HOW TO USE THE TABLE:

Pests		Crops: within the Cucurbitaceae		Crops: outside Cucurbitaceae	
1	2	3	4	5	6
Pest species	Pest group name	Indicator crops	Extrapolation to other crops	Data from these crops can support the indicator crops (reduced data or no data *)	Extrapolation to crops (reduced or no data*)
<i>Delia platura</i> HYLEPL	Root and soil flies	Melon CUMME or Cucumber CUMSC	All crops within the crop group	Field bean VICFX , potato SOLTU, Soybean GLXMA, <i>Phaseolus</i> sp. PHSSS, spinach SPQOL, asparagus ASPOF, Allium vegetables	<i>Freesia</i> sp. FRESS, Allium vegetables, Asparagus ASPOF

E.g. : In the first row above, in order to support a claim for *Delia platura* on all Cucurbitaceae crops, data can be generated either on cucumber, or melon. The number of trials required on these crops can be reduced if there are existing relevant data for *Delia platura* on field bean or potato or soybean or *Phaseolus* spp. or spinach or asparagus or allium vegetables. Data on *Delia platura* generated on Cucurbitaceae can also be used to support claims on a minor use crop such as Freesia, Allium vegetables or Asparagus, but further additional data may be required. The company may also need to consider and justify the minor use status of the specified crop.

EXTRAPOLATION REGARDING PROTECTED/OUTDOOR SITUATIONS

Please note that where crops may be grown in both protected and field situations, and where significant differences are expected in pest relevance or crop agronomy between indoor and outdoor situations, it is important to generate a proportion of the data on crops grown in both situations to ensure the product has been tested under a suitable range of typical and challenging conditions.

EXTRAPOLATION TABLE for EFFECTIVENESS of INSECTICIDES

► PESTS ON STONE FRUIT

PRNPS peach *Prunus persica* including, PRNPN nectarine *P. persica* var. *nucipersica* (and similar hybrids), PRNAR apricot *P. armeniaca*, PRNDU almond *P. dulcis*, PRNDO plum *P. domestica*, PRNDD damson plum *P. damascene*, PRNDS mirabelle *P. domestica* var. *Syriaca*, PRNDI greengage (reine-claude) *P. domestica* subsp. *italic*, PRNDT bullace *P. domestica* subsp. *Insititia*, PRNSN sloe *P. spinosa*, PRNSC Japanese plum *P. salicina*, PRNAV sweet cherry *P. avium*, PRNCE sour cherry *P. cerasus*

Pests		Crops: within Stone Fruit		Crops: outside Stone Fruit	
1 Pest species	2 Pest group name	3 Indicator crops	4 Extrapolation to other crops	5 Data from these crops can support the indicator crops (reduced data or no data *)	6 Extrapolation to crops (reduced or no data*)
<i>Myzus persicae</i> MYZUPE	Aphids	Peach PRNPS	Nectarine PRNPN, Apricot PRNAR, Almond PRNDU, Plum PRNDO		
<i>M. cerasi</i> MYZUCE		Cherry PRNAV	Sour cherry PRNCE		
<i>Appelia prunicola</i> (= <i>Brachycaudus Prunicola</i>) APPEPR, <i>Hyaloptersus pruni</i> HYALPR desirable	Aphids Extrapolation on <i>Phorodon humuli</i> not possible	Plum PRNDO Almonds PRNDU	Damson PRNDD		
<i>Pseudaulacaspis pentagona</i> PSEAPE, <i>Parthenolecanium corni</i> LECACO, <i>Eulecanium tiliae</i> LECATI	Soft scales <i>Coccidae</i>	Peach PRNPS	Apricot PRNAR, Nectarine PRNPN, Almond PRNDU	Other relevant crops	Kiwi ATIDE Ornamental trees and shrubs
<i>Metcalfa pruinosa</i> METFPR	Leafhoppers	Peach PRNPS	Stone fruit	Apple MABSD, Pear PYUCO	Pome fruit Ornamental trees
<i>Aculus schlehtendali</i> VASASD,	Rust mites	Plum PRNDO	Plum PRNDO		Pome fruit Ornamental trees,

<i>Aculus fockeui</i> VASAFL, <i>Phyllocoptes gracillis</i> ACEIGR					Raspberries RUBID (because of <i>Phyllocoptes gracillis</i>)
<i>Eptrimerus pyri</i> EPITPI, <i>Eriophyes pyri</i> ERPHPI, <i>Eriophyes similis</i> ERPHSI	Gall mites <i>Eriophyidae</i>	Plum PRNDO			Pome fruit Ornamental trees Raspberries RUBID (because of <i>Phyllocoptes gracillis</i>)
<i>Panonychus ulmi</i> METTUL, <i>Tetranychus urticae</i> TETRUR, <i>T. viennensis</i> TETRVI, <i>Bryobia rubricolus</i> BRYORU	Spider mites	Peach PRNPS	Stone fruit	Citrus fruit	Pome fruit Other relevant crops Ornamental trees and shrubs
<i>Adoxophyes orana</i> CAPURE and one of the following: <i>Archips podana</i> (= <i>Cacoecia podana</i>) CACOPO, <i>Archips rosanus</i> (= <i>Cacoecia</i> <i>rosana</i>) CACORO, <i>Pandemis heparana</i> PANDHE, <i>Hedya</i> <i>dimidioalba</i> (= <i>Argyroploce</i> <i>variegana</i> , <i>H. nubiferana</i>) ARGPVA, <i>Clepsia spectrana</i> (= <i>Cacoecia costana</i>) CACOCO, <i>Enarmonia</i> <i>formosana</i> ENARFO, <i>Hedya</i> <i>pruniana</i> (= <i>Argyroploce</i> <i>pruniana</i>) ARGPPR, <i>Spilonota ocellana</i> (= <i>Tmetocera ocellana</i>) TMETOC	<i>Tortricidae</i> (Leaf roller moths)	Peach PRNPS	Stone fruit	Any other pome fruit	Pome fruit Ornamental trees
<i>Cydia funebrana</i> LASPFU	<i>Tortricidae</i> (leaf roller moths)	Plum PRNDO	Damsons PRNDD all subspecies of <i>P.</i> <i>domestica</i>		

<i>Cydia.Molesta</i> LAPSMO		Peach PRNPS	Apricot PRNAR, Plum PRNDO		Pome fruit
Data on any two of: <i>Lyonetia clerkella</i> LYONCL, <i>Phyllonorycter blancardella</i> LITHBL, <i>P. corylifoliella</i> PRYCCO, <i>Leucoptera malifoliella</i> LEUCSC, <i>Stigmella malella</i> NEPTMA	Leaf miners	Any relevant stone fruit	Stone fruit		Pome fruit
<i>Anarsia lineatella</i> ANARLI	Stem borers	Peach PRNPS	Stone fruit		
<i>Orthosia</i> sp. ORTOSP	Noctuids	Cherry PRNAV or Plum PRNDO	Stone fruit	Ornamental shrubs	Pome fruit
<i>Operophtera brumata</i> CHEIBR	Winter moths Geometridae	Cherry PRNAV or Plum PRNDO	Stone fruit	Ornamental shrubs	Pome fruit
<i>Orgyia antiqua</i> ORGYAN	<i>Lymantriidae</i>	Cherry PRNAV or Plum PRNDO	Stone fruit	Ornamental shrubs	Pome fruit
<i>Taxonus glabratus</i> TAXOGL, <i>Hoplocampa</i> sp., HOPLSP, <i>Caliroa cerasi</i> (= <i>Eriocampoides limacine</i>) ERICLI	Sawflies	Any relevant stone fruit	Stone fruit		Pome fruit
<i>Ceratitis capitata</i> CERTCA	Fruit flies	Peach PRNPS	Stone fruit	Citrus fruit	Pome fruit Kiwi* ATIDE Kaki DOSKA Figs FIUCA, Litchi LIHCH, Guava PSIGU, Annona fruits, Mango MNGIN, Papaya CIAPA, Passion fruit PAQUED
<i>Rhagoletis cerasi</i> RHAGCE	Fruit flies	Cherry PRNAV			

<i>Drosophila suzukii</i> DROSSU	Fruit flies	Cherry PRNAV	Stone fruit	Currant and berries, Strawberry FRAAN	Currant and berries, Grapes VITVI Strawberries FRAAN, Figs FIUCA, Persimmon DOSKA, Kiwi ATIDE
<i>Peritelus sphaeroides</i> PERESH, <i>Polydrusus sp</i> POLOSP, <i>Phyllobius sp</i> PLL BSP, <i>Otiorhynchus</i> <i>impressiventris</i> OTIOIM, <i>Byctiscus betulae</i> BYCTBE	Weevils (leaf- eaters)	Any stone fruit	Stone fruit		Pome fruit
<i>Rhynchites coeruleus</i> RNCHCA, <i>R. pauxillus</i> COENPA	Weevils (bud- damaging)	Any stone fruit	Stone fruit		Pome fruit
<i>Rhynchites bacchus</i> RNCHBA, <i>Rhynchites</i> <i>aequatus</i> COENAQ, <i>Furcipes rectirostris</i> ANTHRE	Weevils (fruit damaging)	Any stone fruit	Stone fruit		Pome fruit
<i>Capnodis tenebrionis</i> CAPNTE	<i>Buprestidae</i>	Apricot PRNAR	Peach PRNPS		
<i>Cacopsylla pruni</i> PSYLPR		European Plum PRNDO and Japanese Plum PRNSC	Stone fruit		
<i>Halyomorpha halys</i> HALYHA		Peach PRNPS	Stone fruit		Pome fruit Ornamental trees and shrubs