

**EXTRAPOLATION TABLE for CROP SAFETY of FUNGICIDES and INSECTICIDES**  
**► VEGETABLE BRASSICAS <sup>a</sup>**

Treatment type		Indicator crops	Extrapolation to other crops	Data from other crops (or crop groups) that enables reduced data on the indicator crops (or no data *)	Data on indicator crops that permits extrapolation to other crops (or crop groups) with reduced data (or no data *)
Type of application	Timing	Crop	Crop	Crop	Crop
Seed treatment <sup>b</sup>		Cauliflower BRSOB	Leafy and flower head and head and root brassicas	Lettuce LACSA, sensitive ornamentals <sup>c</sup>	
Soil treatment		Any			
Foliar treatment	<i>Before heading</i>	Chinese cabbage BRSCH			
	<i>After heading</i>	Chinese cabbage BRSCH <u>and</u> Broccoli BRSOK <u>or</u> cauliflower BRSOB			

<sup>a</sup> For vegetable brassicas, specific trials with insecticides and fungicides are not essential for foliar treatment. Observations in efficacy or residue trials are usually acceptable. For seed treatment a germination study on the indicator crop is usually necessary.

<sup>b</sup> For seed treatments, indicator crops should include seeds of similar or smaller size.

<sup>c</sup> See the extrapolation table n° 11 (PSD/EU DG SANCO tables) on EPPO website: [http://www.eppo.org/PPPRODUCTS/minor\\_uses/extrapolation\\_tables.pdf](http://www.eppo.org/PPPRODUCTS/minor_uses/extrapolation_tables.pdf)